

**Reference List 2007-2011**

No. 265 - 376

**2007**

- (265) Axelsson CK, Mouridsen HT, Düring M, Moller S. Axillary staging during surgery for breast cancer. *Br J Surg* 2007 Mar;94(3):304-9.
- (266) Dalton SO, Ross L, Düring M, Carlsen K, Mortensen PB, Lynch J, Johansen C. Influence of socioeconomic factors on survival after breast cancer--a nationwide cohort study of women diagnosed with breast cancer in Denmark 1983-1999. *Int J Cancer*. 2007 Dec 1;121(11):2524-31.
- (267) Ejlertsen B, Hojris I, Hansen S, Moholt K, Kristensen B, Mouridsen HT et al. WITHDRAWN: Combined epirubicin and vinorelbine as first-line therapy in metastatic breast cancer: a pilot study performed by the Danish Breast Cancer Cooperative Group. *Breast* 2007 May 29.
- (268) Ejlertsen B, Mouridsen HT, Jensen MB, Andersen J, Cold S, Edlund P et al. Improved outcome from substituting methotrexate with epirubicin: results from a randomised comparison of CMF versus CEF in patients with primary breast cancer. *Eur J Cancer* 2007 March;43(5):877-84.
- (269) Grabau D, Jensen MB, Rank F, Blichert-Toft M. Axillary lymph node micrometastases in invasive breast cancer: national figures on incidence and overall survival. *APMIS* 2007 July;115(7):828-37.
- (270) Groenvold M, Fayers PM, Petersen MA, Sprangers MA, Aaronson NK, Mouridsen HT. Breast cancer patients on adjuvant chemotherapy report a wide range of problems not identified by health-care staff. *Breast Cancer Res Treat* 2007 June;103(2):185-95.
- (271) Groenvold M, Petersen MA, Idler E, Bjorner JB, Fayers PM, Mouridsen HT. Psychological distress and fatigue predicted recurrence and survival in primary breast cancer patients. *Breast Cancer Res Treat* 2007 October;105(2):209-19.
- (272) Gronvold M, Jensen AB. [Patients' experience of breast cancer treatment]. *Ugeskr Laeger* 2007 September 10;169(37):3114-7.
- (273) Husted MA, Haugaard K, Soerensen J, Bokmand S, Friis E, Holtveg H et al. Arm morbidity following sentinel lymph node biopsy or axillary lymph node dissection: A study from the Danish Breast Cancer Cooperative Group. *Breast* 2007 October 8.
- (274) Johansen J, Overgaard J, Overgaard M. Effect of adjuvant systemic treatment on cosmetic outcome and late normal-tissue reactions after breast conservation. *Acta Oncol* 2007;46(4):525-33.
- (275) Offersen BV, Riisbro R, Knoop A, Brunner N, Overgaard J. Lack of association between level of Plasminogen Activator Inhibitor-1 and estimates of tumor angiogenesis in early breast cancer. *Acta Oncol* 2007;46(6):782-91.
- (276) Overgaard M, Nielsen HM, Overgaard J. Is the benefit of postmastectomy irradiation limited to patients with four or more positive nodes, as recommended in international consensus reports? A subgroup analysis of the DBCG 82 b&c randomized trials. *Radiother Oncol* 2007 March;82(3):247-53.

(277) Peuckmann V, Ekholm O, Rasmussen NK, Moller S, Groenvold M, Christiansen P et al. Health-related quality of life in long-term breast cancer survivors: nationwide survey in Denmark. *Breast Cancer Res Treat* 2007 July;104(1):39-46.

(278) Talman ML, Jensen MB, Rank F. Invasive lobular breast cancer. Prognostic significance of histological malignancy grading. *Acta Oncol* 2007;46(6):803-9.

## 2008

(279) Ahern TP, Larsson H, Garne JP, Cronin-Fenton DP, Sorensen HT, Lash TL. Trends in breast-conserving surgery in Denmark, 1982-2002. *Eur J Epidemiol* 2008;23(2):109-14.

(280) Alsner J, Jensen V, Kyndi M, Offersen BV, Vu P, Borresen-Dale AL et al. A comparison between p53 accumulation determined by immunohistochemistry and TP53 mutations as prognostic variables in tumours from breast cancer patients. *Acta Oncol* 2008;47(4):600-7.

(281) Andersen J, Kamby C, Ejlersen B, Cold S, Ewertz M, Jacobsen EH et al. Tamoxifen for one year versus two years versus 6 months of Tamoxifen and 6 months of megestrol acetate: a randomized comparison in postmenopausal patients with high-risk breast cancer (DBCG 89C). *Acta Oncol* 2008;47(4):718-24.

(282) Andersen SB, Vejborg I, von Euler-Chelpin M. Participation behaviour following a false positive test in the Copenhagen mammography screening programme. *Acta Oncol* 2008;47(4):550-5.

(283) Andersson M, Jensen MB, Engholm G, Henrik SH. Risk of second primary cancer among patients with early operable breast cancer registered or randomised in Danish Breast Cancer cooperative Group (DBCG) protocols of the 77, 82 and 89 programmes during 1977-2001. *Acta Oncol* 2008;47(4):755-64.

(284) Axelsson CK, Jonsson PE. Sentinel lymph node biopsy in operations for recurrent breast cancer. *Eur J Surg Oncol* 2008 Jun;34(6):626-30.

(285) Bjarnason NH, Hitz M, Jorgensen NR, Vestergaard P. Adverse bone effects during pharmacological breast cancer therapy. *Acta Oncol* 2008;47(4):747-54.

(286) Blichert-Toft M, Nielsen M, Durning M, Moller S, Rank F, Overgaard M et al. Long-term results of breast conserving surgery vs. mastectomy for early stage invasive breast cancer: 20-year follow-up of the Danish randomized DBCG-82TM protocol. *Acta Oncol* 2008;47(4):672-81.

(287) Blichert-Toft M, Christiansen P, Mouridsen HT. Danish Breast Cancer Cooperative Group--DBCG: History, organization, and status of scientific achievements at 30-year anniversary. *Acta Oncol* 2008;47(4):497-505.

(288) Christensen S, Zachariae R, Jensen AB, Vaeth M, Moller S, Ravnsbaek J et al. Prevalence and risk of depressive symptoms 3-4 months post-surgery in a nationwide cohort study of Danish women treated for early stage breast-cancer. *Breast Cancer Res Treat* 2008 February 16.

(289) Christiansen P, Al-Suliman N, Bjerre K, Moller S. Recurrence pattern and prognosis in low-risk breast cancer patients--data from the DBCG 89-A programme. *Acta Oncol* 2008;47(4):691-703.

(290) Christiansen P, Friis E, Balslev E, Jensen D, Moller S. Sentinel node biopsy in breast cancer: five years experience from Denmark. *Acta Oncol* 2008;47(4):561-8.

- (291) Ejlertsen B, Mouridsen HT, Jensen MB. Adjuvant cyclophosphamide, methotrexate, and fluorouracil in premenopausal patients with node-positive breast cancer: indirect comparison of dose and schedule in DBCG trials 77, 82, and 89. *Acta Oncol* 2008;47(4):662-71.
- (292) Ejlertsen B, Jensen MB, Mouridsen HT, Andersen J, Cold S, Jakobsen E et al. DBCG trial 89B comparing adjuvant CMF and ovarian ablation: similar outcome for eligible but non-enrolled and randomized breast cancer patients. *Acta Oncol* 2008;47(4):709-17.
- (293) Ewertz M, Kempel MM, Duing M, Jensen MB, Andersson M, Christiansen P et al. Breast conserving treatment in Denmark, 1989-1998. A nationwide population-based study of the Danish Breast Cancer Co-operative Group. *Acta Oncol* 2008;47(4):682-90.
- (294) Friis E, Galatius H, Garne JP. Organized nation-wide implementation of sentinel lymph node biopsy in Denmark. *Acta Oncol* 2008;47(4):556-60.
- (295) Henriksen KL, Sonne-Hansen K, Kirkegaard T, Frogne T, Lykkesfeldt AE. Development of new predictive markers for endocrine therapy and resistance in breast cancer. *Acta Oncol* 2008;47(4):795-801.
- (296) Hernandez RK, Sorensen HT, Jacobsen J, Pedersen L, Lash TL. Tamoxifen treatment in Danish breast cancer patients and 5-year risk of arterial atherosclerotic events: a null association. *Cancer Epidemiol Biomarkers Prev* 2008 Sep;17(9):2509-11.
- (297) Holmich LR, Duing M, Henriksen TF, Krag C, Tange UB, Kjoller K, et al. Delayed breast reconstruction with implants after invasive breast cancer does not impair prognosis. *Ann Plast Surg* 2008 Jul;61(1):11-8.
- (298) Husted MA, Haugaard K, Soerensen J, Bokmand S, Friis E, Holtveg H, et al. Arm morbidity following sentinel lymph node biopsy or axillary lymph node dissection: a study from the Danish Breast Cancer Cooperative Group. *Breast* 2008 Apr;17(2):138-47.
- (299) Jensen AR, Madsen AH, Overgaard J. Trends in breast cancer during three decades in Denmark: stage at diagnosis, surgical management and survival. *Acta Oncol* 2008;47(4):537-44.
- (300) Joergensen LE, Gunnarsdottir KA, Lanng C, Moeller S, Rasmussen BB. Multifocality as a prognostic factor in breast cancer patients registered in Danish Breast Cancer Cooperative Group (DBCG) 1996-2001. *Breast* 2008 August 8.
- (301) Johansen H, Kaae S, Jensen MB, Mouridsen HT. Extended radical mastectomy versus simple mastectomy followed by radiotherapy in primary breast cancer. A fifty-year follow-up to the Copenhagen Breast Cancer randomised study. *Acta Oncol* 2008;47(4):633-8.
- (302) Kiaer HW, Laenkholm AV, Nielsen BB, Bjerre KD. Classical pathological variables recorded in the Danish Breast Cancer Cooperative Group's register 1978-2006. *Acta Oncol* 2008;47(4):778-83.
- (303) Kristensen B, Ejlertsen B, Mouridsen HT, Jensen MB, Andersen J, Bjerregaard B et al. Bisphosphonate treatment in primary breast cancer: results from a randomised comparison of oral pamidronate versus no pamidronate in patients with primary breast cancer. *Acta Oncol* 2008;47(4):740-6.
- (304) Kroman N. Timing of breast cancer surgery in relation to the menstrual cycle--the rise and fall of a hypothesis. *Acta Oncol* 2008;47(4):576-9.

- (305) Kroman N, Jensen MB, Wohlfahrt J, Ejlersen B. Pregnancy after treatment of breast cancer--a population-based study on behalf of Danish Breast Cancer Cooperative Group. *Acta Oncol* 2008;47(4):545-9.
- (306) Kyndi M, Sorensen FB, Knudsen H, Overgaard M, Nielsen HM, Andersen J et al. Tissue microarrays compared with whole sections and biochemical analyses. A subgroup analysis of DBCG 82 b&c. *Acta Oncol* 2008;47(4):591-9.
- (307) Kyndi M, Sorensen FB, Knudsen H, Alsner J, Overgaard M, Nielsen HM et al. Impact of BCL2 and p53 on postmastectomy radiotherapy response in high-risk breast cancer. A subgroup analysis of DBCG82 b&c. *Acta Oncol* 2008;47(4):608-17.
- (308) Kyndi M, Overgaard M, Nielsen HM, Sorensen FB, Knudsen H, Overgaard J. High local recurrence risk is not associated with large survival reduction after postmastectomy radiotherapy in high-risk breast cancer: A subgroup analysis of DBCG 82 b&c. *Radiother Oncol* 2008 May 7.
- (309) Kyndi M. Biological markers and response to adjuvant radiotherapy i high-risk breast cancer Faculty of health sciences, University of Aarhus; 2008.
- (310) Kyndi M, Sorensen FB, Knudsen H, Jan A, Overgaard M, Nielsen HM et al. Carbonic anhydrase IX and response to postmastectomy radiotherapy in high-risk breast cancer: a subgroup analysis of the DBCG82 b and c trials. *Breast Cancer Res* 2008 March 20;10(2):R24.
- (311) Kyndi M, Sorensen FB, Knudsen H, Overgaard M, Nielsen HM, Overgaard J. Estrogen receptor, progesterone receptor, HER-2, and response to postmastectomy radiotherapy in high-risk breast cancer: the Danish Breast Cancer Cooperative Group. *J Clin Oncol* 2008 March 20;26(9):1419-26.
- (312) Laenkholm AV, Jensen MB, Kroman N, Rank F. Breast cancer in situ. From pre-malignant lesion of uncertain significance to well-defined non-invasive malignant lesion. The Danish Breast Cancer Cooperative Group Register 1977-2007 revisited. *Acta Oncol* 2008;47(4):765-71.
- (313) Langkjer ST, Ejlersen B, Mouridsen H, Andersen J, Nielsen MM, Moller KA et al. Vinorelbine as first-line or second-line therapy for advanced breast cancer: a Phase I-II trial by the Danish Breast Cancer Co-operative Group. *Acta Oncol* 2008;47(4):735-9.
- (314) Lauridsen MC, Overgaard M, Overgaard J, Hessov IB, Christiansen P. Shoulder disability and late symptoms following surgery for early breast cancer. *Acta Oncol* 2008;47(4):569-75.
- (315) Madsen AH, Jensen AR, Christiansen P, Garne JP, Cold S, Ewertz M et al. Does the introduction of sentinel node biopsy increase the number of node positive patients with early breast cancer? A population based study from the Danish Breast Cancer Cooperative Group. *Acta Oncol* 2008;47(2):239-47.
- (316) Madsen AH, Haugaard K, Soerensen J, Bokmand S, Friis E, Holtveg H et al. Arm morbidity following sentinel lymph node biopsy or axillary lymph node dissection: A study from the Danish Breast Cancer Cooperative Group. *Breast* 2008 April;17(2):138-47.
- (317) Moller S, Jensen MB, Ejlersen B, Bjerre KD, Larsen M, Hansen HB et al. The clinical database and the treatment guidelines of the Danish Breast Cancer Cooperative Group (DBCG); its 30-years experience and future promise. *Acta Oncol* 2008;47(4):506-24.
- (318) Mouridsen HT, Bjerre KD, Christiansen P, Jensen MB, Moller S. Improvement of prognosis in breast cancer in Denmark 1977-2006, based on the nationwide reporting to the DBCG Registry. *Acta Oncol* 2008;47(4):525-36.

- (319) Møller S, Jensen MB, Ejlersen B, Bjerre KD, Larsen M, Hansen HB et al. The clinical database of the Danish Breast Cancer Cooperative Group (DBCG); its 30-years experience and future promise. *Acta Oncol* 2008.
- (320) Nielsen KV, Ejlersen B, Moller S, Jorgensen JT, Knoop A, Knudsen H et al. The value of TOP2A gene copy number variation as a biomarker in breast cancer: Update of DBCG trial 89D. *Acta Oncol* 2008;47(4):725-34.
- (321) Offersen BV, Alsner J, Ege OK, Riisbro R, Brunner N, Sorensen FB et al. A comparison among HER2, TP53, PAI-1, angiogenesis, and proliferation activity as prognostic variables in tumours from 408 patients diagnosed with early breast cancer. *Acta Oncol* 2008;47(4):618-32.
- (322) Overgaard J. DBCG: the Danish Breast Cancer Cooperative Group--a 30-year struggle for better breast cancer treatment in Denmark. *Acta Oncol* 2008;47(4):491-6.
- (323) Overgaard M, Christensen JJ. Postoperative radiotherapy in DBCG during 30 years. Techniques, indications and clinical radiobiological experience. *Acta Oncol* 2008;47(4):639-53.
- (324) Rasmussen BB, Andersson M, Christensen IJ, Moller S. Evaluation of and quality assurance in HER2 analysis in breast carcinomas from patients registered in Danish Breast Cancer Group (DBCG) in the period of 2002-2006. A nationwide study including correlation between HER-2 status and other prognostic variables. *Acta Oncol* 2008;47(4):784-8.
- (325) Roslind A, Knoop AS, Jensen MB, Johansen JS, Nielsen DL, Price PA, Balslev E. YKL-40 protein expression is not a prognostic marker in patients with primary breast cancer. *Breast Cancer Res Treat* 2008 Nov;112(2):275-85
- (326) Stralman K, Mollerup CL, Kristoffersen US, Elberg JJ. Long-term outcome after mastectomy with immediate breast reconstruction. *Acta Oncol* 2008;47(4):704-8.
- (327) Talman ML, Rasmussen BB, Andersen J, Christensen IJ. Estrogen Receptor analyses in the Danish Breast Cancer Cooperative Group. History, methods, prognosis and clinical implications. *Acta Oncol* 2008;47(4):789-94.
- (328) Thomassen M, Hansen TV, Borg A, Lianee HT, Wikman F, Pedersen IS et al. BRCA1 and BRCA2 mutations in Danish families with hereditary breast and/or ovarian cancer. *Acta Oncol* 2008;47(4):772-7.
- (329) Thomsen MS, Berg M, Nielsen HM, Pedersen AN, Overgaard M, Ewertz M et al. Post-mastectomy radiotherapy in Denmark: from 2D to 3D treatment planning guidelines of The Danish Breast Cancer Cooperative Group. *Acta Oncol* 2008;47(4):654-61.
- (330) Würtz SØ, Møller S, Mouridsen H, Hertel PB, Friis E, Brüner N. Plasma and serum levels of tissue inhibitor of metalloproteinases-1 are associated with prognosis in node-negative breast cancer: a prospective study. *Mol Cell Proteomics*. 2008 Feb; 7(2): 424-30.
- (331) Wurtz SO, Schrohl AS, Mouridsen H, Brunner N. TIMP-1 as a tumor marker in breast cancer--an update. *Acta Oncol* 2008;47(4):580-90.

## 2009

- (332) Ahern TP, Pedersen L, Cronin-Fenton DP, Sorensen HT, Lash TL. No increase in breast cancer recurrence with concurrent use of tamoxifen and some CYP2D6-inhibiting medications. *Cancer Epidemiol Biomarkers Prev* 2009 Sep;18(9):2562-4.

- (333) Axelsson CK, Durning M, Christiansen PM, Wamberg PA, Soe KL, Moller S. Impact on regional recurrence and survival of axillary surgery in women with node-negative primary breast cancer. *Br J Surg* 2009 Jan;96(1):40-6.
- (334) Christensen S, Zachariae R, Jensen AB, Vaeth M, Moller S, Ravnsbaek J, et al. Prevalence and risk of depressive symptoms 3-4 months post-surgery in a nationwide cohort study of Danish women treated for early stage breast-cancer. *Breast Cancer Res Treat* 2009 Jan;113(2):339-55.
- (335) Ejlertsen B, Jensen MB, Rank F, Rasmussen BB, Christiansen P, Kroman N, et al. Population-based study of peritumoral lymphovascular invasion and outcome among patients with operable breast cancer. *J Natl Cancer Inst* 2009 May 20;101(10):729-35.
- (336) Gärtner R, Jensen MB, Nielsen J, Ewertz M, Kroman N, Kehlet H. Prevalence of and factors associated with persistent pain following breast cancer surgery. *JAMA*. 2009 Nov 11;302(18):1985-92.
- (337) Henriksen KL, Rasmussen BB, Lykkesfeldt AE, Møller S, Ejlertsen B, Mouridsen HT. An ER activity profile including ER, PR, Bcl-2 and IGF-IR may have potential as selection criterion for letrozole or tamoxifen treatment of patients with advanced breast cancer. *Acta Oncol*. 2009; 48(4):522-31.
- (338) Hernandez RK, Sorensen HT, Pedersen L, Jacobsen J, Lash TL. Tamoxifen treatment and risk of deep venous thrombosis and pulmonary embolism: a Danish population-based cohort study. *Cancer* 2009 Oct 1;115(19):4442-9.
- (339) Kyndi M, Overgaard M, Nielsen HM, Sorensen FB, Knudsen H, Overgaard J. High local recurrence risk is not associated with large survival reduction after postmastectomy radiotherapy in high-risk breast cancer: a subgroup analysis of DBCG 82 b&c. *Radiother Oncol* 2009 Jan;90(1):74-9.
- (340) Lykkesfeldt AE, Henriksen KL, Rasmussen BB, Sasano H, Evans DB, Møller S, Ejlertsen B, Mouridsen HT. In situ aromatase expression in primary tumor is associated with estrogen receptor expression but is not predictive of response to endocrine therapy in advanced breast cancer. *BMC Cancer*. 2009 Jun 16; 9: 185.
- (341) Olsen AH, Bihmann K, Jensen MB, Vejborg I, Lyng E. Breast density and outcome of mammography screening: a cohort study. *Br J Cancer* 2009 Apr 7;100(7):1205-8.
- (342) Peuckmann V, Ekholm O, Rasmussen NK, Groenvold M, Christiansen P, Moller S, et al. Chronic pain and other sequelae in long-term breast cancer survivors: nationwide survey in Denmark. *Eur J Pain* 2009 May;13(5):478-85.
- (343) Peuckmann V, Ekholm O, Sjogren P, Rasmussen NK, Christiansen P, Moller S, et al. Health care utilisation and characteristics of long-term breast cancer survivors: nationwide survey in Denmark. *Eur J Cancer* 2009 Mar;45(4):625-33.
- (344) Willemoe GL, Hertel PB, Bartels A, Jensen MB, Balslev E, Rasmussen BB, et al. Lack of TIMP-1 tumour cell immunoreactivity predicts effect of adjuvant anthracycline-based chemotherapy in patients (n=647) with primary breast cancer. A Danish Breast Cancer Cooperative Group Study. *Eur J Cancer* 2009 Sep;45(14):2528-36.

## 2010

- (345) Blamey RW, Hornmark-Stenstam B, Ball G, Blichert-Toft M, Cataliotti L, Fourquet A, Gee J, Holli K, Jakesz R, Kerin M, Mansel R, Nicholson R, Pienkowski T, Pinder S, Sundquist M, van de

- Vijver M, Ellis I. ONCOPOOL - a European database for 16,944 cases of breast cancer. *Eur J Cancer* 2010 Jan;46(1):56-71.
- (346) Ejlertsen B, Jensen MB, Nielsen KV, Balslev E, Rasmussen BB, Willemoe GL, Hertel PB, Knoop AS, Mouridsen HT, Brüner N. HER2, TOP2A, and TIMP-1 and Responsiveness to Adjuvant Anthracycline-Containing Chemotherapy in High-Risk Breast Cancer Patients. *J Clin Oncol* 2010 Feb 20;28(6):984-90.
- (347) Ejlertsen B, Mouridsen HT, Jensen MB, Andersen J, Andersson M, Kamby C, Knoop AS; Danish Breast Cancer Cooperative Group. Cyclophosphamide, methotrexate, and fluorouracil; oral cyclophosphamide; levamisole; or no adjuvant therapy for patients with high-risk, premenopausal breast cancer. *Cancer*. 2010 May 1;116(9):2081-9.
- (348) Gunnarsdóttir KA, Jensen MB, Zahrieh D, Gelber RD, Knoop A, Bonetti M, Mouridsen H, Ejlertsen B. CEF is superior to CMF for tumours with TOP2A aberrations: a Subpopulation Treatment Effect Pattern Plot (STEPP) analysis on Danish Breast Cancer Cooperative Group Study 89D. *Breast Cancer Res Treat*. 2010 Aug;123(1):163-9.
- (349) Gärtner R, Jensen MB, Kronborg L, Ewertz M, Kehlet H, Kroman N. Self-reported arm-lymphedema and functional impairment after breast cancer treatment--a nationwide study of prevalence and associated factors. *Breast*. 2010 Dec;19(6):506-15.
- (350) Mouridsen HT, Lønning P, Beckmann MW, Blackwell K, Doughty J, Gligorov J, Llombart-Cussac A, Robidoux A, Thürlimann B, Gnant M. Use of aromatase inhibitors and bisphosphonates as an anticancer therapy in postmenopausal breast cancer. *Anticancer Ther*. 2010; 10 (11): 1825-1836.
- (351) Nielsen KV, Müller S, Møller S, Schønau A, Balslev E, Knoop AS, Ejlertsen B. Aberrations of ERBB2 and TOP2A genes in breast cancer. *Mol Oncol*. 2010 Apr; 4(2): 161-8.
- 2011**
- (352) Andersson M, Lidbrink E, Bjerre K, Wist E, Enevoldsen K, Jensen AB, Karlsson P, Tange UB, Sørensen PG, Møller S, Bergh J, Langkjær ST. Phase III randomized study comparing docetaxel plus trastuzumab with vinorelbine plus trastuzumab as first-line therapy of metastatic or locally advanced human epidermal growth factor receptor 2-positive breast cancer: the HERNATA study. *J Clin Oncol*. 2011 Jan 20; 29(3): 264-71.
- (353) Azim HA Jr, Santoro L, Pavlidis N, Gelber S, Kroman N, Azim H, Peccatori FA. Safety of pregnancy following breast cancer diagnosis: A meta-analysis of 14 studies. *Eur J Cancer* 2011 Jan;47(1):74-83.
- (354) Christiansen P, Bjerre K, Ejlertsen B, Jensen M-B, Rasmussen BB, Lænkholm A-V, Kroman N, Ewertz M, Offersen B, Toftdahl DB, Møller S, Mouridsen HT; on behalf of the Danish Breast Cancer Cooperative Group. Mortality Rates Among Early-Stage Hormone Receptor-Positive Breast Cancer Patients: A Population-Based Cohort Study in Denmark. *J Natl Cancer Inst*. 2011 Aug [Epub ahead of print].
- (355) Damkjær LH, Deltour I, Suppli NP, Christensen J, Kroman NT, Johansen C & Dalton SO. Breast cancer and early retirement: Associations with disease characteristics, treatment, comorbidity, social position and participation in a six-day rehabilitation course in a register-based study in Denmark. *Acta Oncol* 2011; 50: 274–281.
- (356) Darby S, McGale P, Correa C, Taylor C, Arriagada R, Clarke M, Cutter D, Davies C, Ewertz M, Godwin J, Gray R, Pierce L, Whelan T, Wang Y and Peto R; The writing committee on behalf of

- the Early Breast Cancer Trialists' Collaborative Group (EBCTCG). Effect of radiotherapy after breast-conserving surgery on 10-year recurrence and 15-year breast cancer death: meta-analysis of individual patient data for 10 801 women in 17 randomised trials. *Lancet* 2011 Oct [Epub ahead of print].
- (357) Di Leo A, Desmedt C, Bartlett JMS, Piette F, Ejlertsen B, Pritchard KI, Larsimont D, Poole C, Isola J, Earl H, Mouridsen H, O'Malley FP, Cardoso F, Tanner M, Munro A, Twelves Prof C J, Sotiriou C, Shepherd L, Cameron D, Piccart MJ, Buyse M, for the HER2/TOP2A Meta-analysis Study Group. HER2 and TOP2A as predictive markers for anthracycline-containing chemotherapy regimens as adjuvant treatment of breast cancer: a meta-analysis of individual patient data. *Lancet Oncol* 2011 Sept [Epub ahead of print].
- (358) Eckhoff L, Nielsen M, Moeller S, Knoop A. TAXTOX - a retrospective study regarding the side effects of docetaxel given as part of the adjuvant treatment to patients with primary breast cancer in Denmark from 2007 to 2009. *Acta Oncol.* 2011 Oct; 50(7): 1075-82.
- (359) Edlund P, Ahlgren J, Bjerre K, Andersson M, Bergh J, Mouridsen H, Holmberg SB, Bengtsson NO, Jakobsen E, Møller S, Lindman H, Blomqvist C. Dose-tailoring of FEC adjuvant chemotherapy based on leukopenia is feasible and well tolerated. Toxicity and dose intensity in the Scandinavian Breast Group phase 3 adjuvant Trial SBG 2000-1. *Acta Oncol.* 2011 Apr; 50(3): 329-37.
- (360) Ejlertsen B, Aldridge J, Nielsen KV, Regan MM, Henriksen KL, Lykkesfeldt AE, Müller S, Gelber RD, Price KN, Rasmussen BB, Viale G, Mouridsen H; for the Danish Breast Cancer Cooperative Group, the BIG 1-98 Collaborative Group and the International Breast Cancer Study Group. Prognostic and predictive role of ESR1 status for postmenopausal patients with endocrine-responsive early breast cancer in the Danish cohort of the BIG 1-98 trial. *Ann Oncol* 2011 Oct 10. [Epub ahead of print]
- (361) Ewertz M, Jensen MB, Gunnarsdóttir KA, Højris I, Jakobsen EH, Nielsen D, Stenbygaard LE, Tange UB, Cold S. Effect of Obesity on Prognosis After Early-Stage Breast Cancer. *J Clin Oncol* 2011 Jan 1;29(1):25-31.
- (362) Hertel, Pernille Bræmer. TIMP-1 as a predictive marker for chemotherapy in primary breast cancer. University of Copenhagen, Department of Veterinary Disease Biology; PhD Thesis 2011.
- (363) Jensen JD, Knoop A, Lænkholm AV, Grauslund M, Jensen M-B, Santoni-Rugiu E, Andersson M, Ewertz M. 2PIK3CA mutations, PTEN, and pHER2 expression and impact on outcome in HER2-positive early-stage breast cancer patients treated with adjuvant chemotherapy and trastuzumab. *Ann Oncol* 2011 Dec [Epub ahead of print].
- (364) Jørgensen JT, Møller S, Rasmussen BB, Winther H, Schønau A, Knoop A. High concordance between two companion diagnostics tests: a concordance study between the HercepTest and the HER2 FISH pharmDx kit. *Am J Clin Pathol.* 2011 Jul; 136(1):145-51.
- (365) Land LH, Dalton SO, Jensen M-B, Ewertz M. Impact of comorbidity on mortality: a cohort study of 62,591 Danish women diagnosed with early breast cancer, 1990-2008. *Breast Cancer Res Treat.* 2011 Oct [Epub ahead of print].
- (366) McGale P, Darby SC, Hall P, Adolfsson J, Bengtsson N-O, Bennet AM, Fornander T, Gigante B, Jensen M-B, Peto R, Rahimi K, Taylor CW, Ewertz M. Incidence of heart disease in 35,000 women treated with radiotherapy for breast cancer in Denmark and Sweden. *Radiother Oncol* 2011; 100: 167-175.

- (367) Nielsen DL, Bjerre KD, Jakobsen EH, Cold S, Stenbygaard L, Sørensen PG, Kamby C, Møller S, Jørgensen CL, Andersson M. Gemcitabine Plus Docetaxel Versus Docetaxel in Patients With Predominantly Human Epidermal Growth Factor Receptor 2-Negative Locally Advanced or Metastatic Breast Cancer: A Randomized, Phase III Study by the Danish Breast Cancer Cooperative Group. *J Clin Oncol*. 2011 Dec 20; 29(36): 4748-54.
- (368) Nielsen KV, Ejlersen B, Müller S, Møller S, Rasmussen BB, Balslev E, Lænkholm AV, Christiansen P, Mouridsen HT. Amplification of ESR1 may predict resistance to adjuvant tamoxifen in postmenopausal patients with hormone receptor positive breast cancer. *Breast Cancer Res Treat* 2011; 127: 345-355.
- (369) Nielsen KV, Ejlersen B, Møller S, Jensen M-B, Balslev E, Müller S, Knoop A, Mouridsen HT. Lack of independent prognostic and predictive value of centromere 17 copy number changes in breast cancer patients with known HER2 and TOP2A status. *Molecular Oncology* 2011 Dec [Epub ahead of print].
- (370) O'Connor M, Christensen S, Jensen AB, Møller S, Zachariae R. How traumatic is breast cancer? Post-traumatic stress symptoms (PTSS) and risk factors for severe PTSS at 3 and 15 months after surgery in a nationwide cohort of Danish women treated for primary breast cancer. *Br J Cancer*. 2011 Feb 1;104(3): 419-26.
- (371) Pedersen AN, Møller S, Steffensen KD, Haahr V, Jensen M, Kempel MM, Jepsen SL, Madsen EL, Roslind A, Sandberg E, Schöllkopf C, Sørensen PG, Windfeldt KM, Andersson M. Supraclavicular recurrence after early breast cancer: a curable condition? *Breast Cancer Res Treat*. 2011 Feb; 125(3): 815-22.
- (372) Suppli NP, Deltour I, Damkjær LH, Christensen J, Jensen AB, Kroman NT, Johansen C & Dalton SO. Factors associated with the prescription of antidepressive medication to breast cancer patients. *Acta Oncol* 2011; 50: 243–251.
- (373) Taylor CW, Brønnum D, Darby SC, Gagliardi G, Hall P, Jensen M-B, McGale P, Nisbet A, Ewertz M. Cardiac dose estimates from Danish and Swedish breast cancer radiotherapy during 1977–2001. *Radiother Oncol* 2011; 100: 176-183.
- (374) Tvedskov TF, Bartels A, Jensen M-B, Paaschburg B, Kroman N, Balslev E and Brüner N. Evaluating TIMP-1, Ki67, and HER2 as markers for non-sentinel node metastases in breast cancer patients with micrometastases to the sentinel node. *APMIS* 2011 Oct [Epub ahead of print].
- (375) Tvedskov TF, Jensen M-B, Balslev E, Ejlersen B, Kroman N. Stage migration after introduction of sentinel lymph node dissection in breast cancer treatment in Denmark: A nationwide study. *Eur J Cancer* 2011 Apr;47(6):872-878.
- (376) Tvedskov TF, Jensen M-B, Kroman N, Balslev E. Iatrogenic displacement of tumor cells to the sentinel node after surgical excision in primary breast cancer. *Breast Cancer Res Treat*. 2011 Aug [Epub ahead of print].